

Haut- und Weichteilinfektionen: Erysipel, Zellulitis und nekrotisierende Fasziitis

Infections de la peau et des tissus mous: cellulite, érysipèle et fasciite nécrosante

Literatur / Références

1. Phoenix G, Das S, Joshi M. Diagnosis and management of cellulitis. *BMJ* 2012;345:e4955.
2. Edelsberg J, Taneja C, Zervos M, Haque N, Moore C, Reyes K, et al. *Emerg Infect Dis* 2009;15:1516–8.
3. Snodgrass WR, Anderson T. Prontosil in the treatment of erysipelas. A controlled series of 312 cases. *Br Med J* 1937;2(3933):101–4.
4. Kilburn SA, Featherstone P, Higgins B, Brindle R. Interventions for cellulitis and erysipelas. *Cochrane Database Syst Rev*. 2010 Jun 16;(6):CD004299. doi: 10.1002/14651858.CD004299.pub2
5. Chira S, Miller LG. *Staphylococcus aureus* is the most common identified cause of cellulitis: a systematic review. *Epidemiol Infect* 2010;138:313–7.
6. Stevens DL, Bisno AL, Chambers HF, Everett ED, Dellinger P, Goldstein EJC et al. Practice guidelines for the diagnosis and management of skin and soft-tissue infections. *Clin Infect Dis* 2005;41:1373–406.
7. Gunderson CG. Cellulitis: definition, etiology and clinical features. *Am J Med* 2011;124:1113–22.
8. Sultan HJ, Boyle AA, Sheppard N. Necrotising fasciitis. *BMJ* 2012;345:e4274.
9. Otter JA, French GL. Molecular epidemiology of community-associated methicillin-resistant *Staphylococcus aureus* in Europe. *Lancet Infect Dis* 2010;10:227–39.
10. Overesch G, Büttner S, Rossano A, Perreten V. The increase of methicillin-resistant *Staphylococcus aureus* (MRSA) and the presence of an unusual sequence type ST49 in slaughter pigs in Switzerland. *BMC Vet Res* 2011;7:30.
11. Valsesia G, Rossi M, Bertschy S, Pfyffer GE. Emergence of SCCmec type IV and SCCmec type V methicillin-resistant *Staphylococcus aureus* containing the Panton-Valentine leukocidin genes in a large academic teaching hospital in central Switzerland: external invaders or persisting circulators? *J Clin Microbiol* 2010;48:720–7.
12. Gottlieb T, Atkins BL, Shaw DR. Soft tissue, bone and joint infections. *Med J Aust* 2002;176:609–15.
13. Eron LJ. In the clinic: cellulitis and soft-tissue infections. *Ann Intern Med* 2009;ITC1–16.
14. Tayal VS, Hasan N, Norton J, Tomaszewski CA. The effect of soft-tissue ultrasound on the management of cellulitis in the emergency department. *Acad Emerg Med* 2006;13:384–8.
15. Liu C, Bayer A, Cosgrove SE, Daum RS, Fridkin SK, Gorwitz RJ et al. Clinical practice guidelines by the Infectious Diseases Society of America for the treatment of methicillin-resistant *Staphylococcus aureus* infections in adults and children. *Clin Infect Dis* 2011;52:1–38.
16. Hepburn MJ, Dooley DP, Skidmore PJ, Ellis MW, Starnes WF, Hasewinkle WC. Comparison of short-course (5 days) and standard (10 days) treatment for uncomplicated cellulitis. *Arch Intern Med* 2004;164:1669–74.
17. Thomas KS, Crook AM, Nunn AJ, Foster KA, Mason JM, Chalmers JR, et al. Penicillin to prevent recurrent leg cellulitis. *N Engl J Med* 2013;368:1695–703.
18. Thomas K, Crook A, Foster K, Mason J, Chalmers J, Bourke J, et al. Prophylactic antibiotics for the prevention of cellulitis (erysipelas) of the leg: results of the U.K. Dermatology Clinical Trials Network's PATCH II trial. *Br J Dermatol* 2012;166:169–78.
19. Sultan HY, Boyle AA, Sheppard N. Necrotising fasciitis. *BMJ* 2012;345:e4274.
20. Ustin JS, Malangoni MA. Necrotizing soft-tissue infections. *Crit Care Med* 2011;39:2156–62.
21. Puvanendran R, Chan Meng Huey J, Pasupathy S. Necrotizing fasciitis. *Can Fam Physician* 2009;55:981–7.
22. Wall DB, Klein SR, Black S, de Virgilio C. A simple model to help distinguish necrotizing fasciitis from nonnecrotizing soft tissue infection. *J Am Coll Surg* 2000;191:227–31.
23. Wong CH, Khin LW, Heng KS, Tan KC, Low CO. The LRINEC (Laboratory Risk Indicator for Necrotizing Fasciitis) score: a tool for distinguishing necrotizing fasciitis from other soft tissue infections. *Crit Care Med* 2004;32:1535–41.
24. Swain RA, Hatcher JC, Azadian BS, Soni N, De Souza B. A five-year review of necrotising fasciitis in a tertiary referral unit. *Ann R Coll Surg Engl* 2013;95:57–60.