

Hat die klinische 7T-MRI-Bildgebung eine Zukunft?

L'IRM à 7 Tesla a-t-elle un avenir en pratique clinique?

Literatur / Références

1. Kangarlu A, Shellock FG. Aneurysm clips: evaluation of magnetic field interactions with an 8.0 T MR system. *Journal of magnetic resonance imaging: JMRI*. Jul 2000;12(1):107–111.
2. Christoforidis GA, Grecula JC, Newton HB, et al. Visualization of microvasculature in glioblastoma multiforme with 8–T high-spatial-resolution MR imaging. *AJNR. American journal of neuroradiology*. Oct 2002;23(9):1553–1556.
3. Abduljalil AM, Schmalbrock P, Novak V, Chakeres DW. Enhanced gray and white matter contrast of phase susceptibility-weighted images in ultra-high-field magnetic resonance imaging. *Journal of magnetic resonance imaging: JMRI*. Sep 2003;18(3):284–290.
4. Dashner RA, Chakeres DW, Kangarlu A, Schmalbrock P, Christoforidis GA, DePhilip RM. MR imaging visualization of the cerebral microvasculature: a comparison of live and postmortem studies at 8 T. *AJNR. American journal of neuroradiology*. Oct 2003;24(9):1881–1884.
5. Christoforidis GA, Kangarlu A, Abduljalil AM, et al. Susceptibility-based imaging of glioblastoma microvasculature at 8 T: correlation of MR imaging and postmortem pathology. *AJNR. American journal of neuroradiology*. May 2004;25(5):756–760.
6. Chakeres DW, Whitaker CD, Dashner RA, et al. High-resolution 8 Tesla imaging of the formalin-fixed normal human hippocampus. *Clin Anat*. Mar 2005;18(2):88–91.
7. Ibrahim TS, Mitchell C, Schmalbrock P, Lee R, Chakeres DW. Electromagnetic perspective on the operation of RF coils at 1.5–11.7 Tesla. *Magnetic resonance in medicine: official journal of the Society of Magnetic Resonance in Medicine / Society of Magnetic Resonance in Medicine*. Sep 2005;54(3):683–690.
8. Richdale K, Wassenaar P, Teal Bluestein K, et al. 7 Tesla MR imaging of the human eye in vivo. *Journal of magnetic resonance imaging: JMRI*. Nov 2009;30(5):924–932.
9. Choi S, Cunningham DT, Aguila F, et al. DTI at 7 and 3 T: systematic comparison of SNR and its influence on quantitative metrics. *Magnetic resonance imaging*. Jul 2011;29(6):739–751.
10. Pitt D, Boster A, Pei W, et al. Imaging cortical lesions in multiple sclerosis with ultra-high-field magnetic resonance imaging. *Archives of neurology*. Jul 2010;67(7):812–818.
11. Bluestein KT, Pitt D, Sammet S, et al. Detecting cortical lesions in multiple sclerosis at 7 T using white matter signal attenuation. *Magnetic resonance imaging*. Sep 2012;30(7):907–915.
12. Chang LS, Jacob A, Lorenz M, et al. Growth of benign and malignant schwannoma xenografts in severe combined immunodeficiency mice. *The Laryngoscope*. Nov 2006;116(11):2018–2026.
13. Yuh WT, Christoforidis GA, Koch RM, et al. Clinical magnetic resonance imaging of brain tumors at ultrahigh field: a state-of-the-art review. *Topics in magnetic resonance imaging: TMRI*. Apr 2006;17(2):53–61.
14. Christoforidis JB, Wassenaar PA, Christoforidis GA, Ho VY, Knopp MV, Schmalbrock PM. Retrobulbar vasculature using 7–T magnetic resonance imaging with dedicated eye surface coil. *Graefes's archive for clinical and experimental ophthalmology = Albrecht von Graefes Archiv für klinische und experimentelle Ophthalmologie*. Jan 2013;251(1):271–277.
15. Mehta V, Pei W, Yang G, et al. Iron is a sensitive biomarker for inflammation in multiple sclerosis lesions. *PLoS one*. 2013;8(3):e57573.
16. Apprich S, Trattinig S, Welsch GH, et al. Assessment of articular cartilage repair tissue after matrix-associated autologous chondrocyte transplantation or the microfracture technique in the ankle joint using diffusion-weighted imaging at 3 Tesla. *Osteoarthritis and cartilage / OARS, Osteoarthritis Research Society*. Jul 2012;20(7):703–711.
17. Juras V, Welsch G, Bar P, Kronnerwetter C, Fujita H, Trattinig S. Comparison of 3T and 7T MRI clinical sequences for ankle imaging. *European journal of radiology*. Aug 2012;81(8):1846–1850.
18. Kraff O, Theysohn JM, Maderwald S, et al. MRI of the knee at 7.0 Tesla. *RoFo: Fortschritte auf dem Gebiete der Röntgenstrahlen und der Nuklearmedizin*. Dec 2007;179(12):1231–1235.
19. Trattinig S, Zbyn S, Schmitt B, et al. Advanced MR methods at ultra-high field (7 Tesla) for clinical musculoskeletal applications. *European radiology*. Nov 2012;22(11):2338–2346.
20. Bogner W, Chmelik M, Schmid AI, Moser E, Trattinig S, Gruber S. Assessment of (31)P relaxation times in the human calf muscle: a comparison between 3 T and 7 T in vivo. *Magnetic resonance in medicine: official journal of the Society of Magnetic Resonance in Medicine / Society of Magnetic Resonance in Medicine*. Sep 2009;62(3):574–582.
21. Juras V, Zbyn S, Pressl C, et al. Sodium MR imaging of Achilles tendinopathy at 7 T: preliminary results. *Radiology*. Jan 2012;262(1):199–205.

22. Madelin G, Jerschow A, Regatte RR. Sodium relaxation times in the knee joint in vivo at 7T. *NMR in biomedicine*. Apr 2012;25(4):530–537.
23. Lanzman RS, Kropil P, Schmitt P, et al. Nonenhanced free-breathing ECG-gated steady-state free precession 3D MR angiography of the renal arteries: comparison between 1.5 T and 3 T. *AJR. American journal of roentgenology*. Mar 2010;194(3):794–798.
24. Heverhagen JT, Bourekas E, Sammet S, Knopp MV, Schmalbrock P. Time-of-flight magnetic resonance angiography at 7 Tesla. *Investigative radiology*. Aug 2008;43(8):568–573.
25. Stamm AC, Wright CL, Knopp MV, Schmalbrock P, Heverhagen JT. Phase contrast and time-of-flight magnetic resonance angiography of the intracerebral arteries at 1.5, 3 and 7 T. *Magnetic resonance imaging*. May 2013;31(4):545–549.
26. Kraff O, Bitz AK, Breyer T, et al. A transmit/receive radiofrequency array for imaging the carotid arteries at 7 Tesla: coil design and first in vivo results. *Investigative radiology*. Apr 2011;46(4):246–254.
27. Fischer A, Maderwald S, Orzada S, et al. Nonenhanced Magnetic Resonance Angiography of the Lower Extremity Vessels at 7 Tesla: Initial Experience. *Investigative radiology*. Mar 13 2013.
28. van Elderen SG, Versluis MJ, Webb AG, et al. Initial results on in vivo human coronary MR angiography at 7 T. *Magnetic resonance in medicine: official journal of the Society of Magnetic Resonance in Medicine / Society of Magnetic Resonance in Medicine*. Dec 2009;62(6):1379–1384.
29. Metzger GJ, Auerbach EJ, Akgun C, et al. Dynamically applied B1+ shimming solutions for non-contrast enhanced renal angiography at 7.0 Tesla. *Magnetic resonance in medicine: official journal of the Society of Magnetic Resonance in Medicine / Society of Magnetic Resonance in Medicine*. Jan 2013;69(1):114–126.