

## Abklärung und Behandlung bei PAVK:

### Aspekte der ambulanten Medizin / Aspekte der Spitalmedizin

## Diagnostic et traitement de l'AOMI:

### aspects de la médecine ambulatoire / aspects de la médecine hospitalière

## Literatur / Références

1. A randomised, blinded, trial of clopidogrel versus aspirin in patients at risk of ischaemic events (CAPRIE). CAPRIE Steering Committee. Lancet. 1996;348(9038):1329–39.
2. Adam DJ, Beard JD, Cleveland T, Bell J, Bradbury AW, Forbes JF, et al. Bypass versus angioplasty in severe ischaemia of the leg (BASIL): multicentre, randomised controlled trial. Lancet. 2005;366(9501):1925–34.
3. Alonso-Coello P, Bellmunt S, McGorrian C, Anand SS, Guzman R, Criqui MH, et al. Antithrombotic therapy in peripheral artery disease: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. Chest. 2012;141(2 Suppl):e669S–90S.
4. Anand S, Yusuf S, Xie C, Pogue J, Eikelboom J, Budaj A, et al. Oral anticoagulant and antiplatelet therapy and peripheral arterial disease. N Engl J Med. 2007;357(3):217–27.
5. Anderson JL, Halperin JL, Albert NM, Bozkurt B, Brindis RG, Curtis LH, et al. Management of Patients With Peripheral Artery Disease (Compilation of 2005 and 2011 ACCF/AHA Guideline Recommendations): A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. Circulation. 2013;127(13):1425–43.
6. Antolovic D, Rakow A, Contin P, Ulrich A, Rahbari NN, Buchler MW, et al. A randomised controlled pilot trial to evaluate and optimize the use of anti-platelet agents in the perioperative management in patients undergoing general and abdominal surgery--the APAP trial (ISRCTN45810007). Langenbecks Arch Surg. 2012;397(2):297–306.
7. Apelqvist J, Elgzyri T, Larsson J, Londahl M, Nyberg P, Thorne J. Factors related to outcome of neuroischemic/ischemic foot ulcer in diabetic patients. J Vasc Surg. 2011;53(6):1582–8 e2.
8. Aronow WS, Ahn C. Prevalence of coexistence of coronary artery disease, peripheral arterial disease, and atherothrombotic brain infarction in men and women > or = 62 years of age. Am J Cardiol. 1994;74(1):64–5.
9. Arozullah AM, Daley J, Henderson WG, Khuri SF. Multifactorial risk index for predicting postoperative respiratory failure in men after major noncardiac surgery. The National Veterans Administration Surgical Quality Improvement Program. Ann Surg. 2000;232(2):242–53.
10. Axixa B, Fishwick G, Bolia A, Thompson MM, London NJ, Bell PR, et al. Complications following peripheral angioplasty. Ann R Coll Surg Engl. 2002;84(1):39–42.
11. Bargellini I, Piaggesi A, Cicorelli A, Rizzo L, Cervelli R, Iacopi E, et al. Predictive value of angiographic scores for the integrated management of the ischemic diabetic foot. J Vasc Surg. 2013.
12. Becattini C, Agnelli G, Schenone A, Eichinger S, Bucherini E, Silingardi M, et al. Aspirin for preventing the recurrence of venous thromboembolism. N Engl J Med. 2012;366(21):1959–67.
13. Berger JS, Hiatt WR. Medical therapy in peripheral artery disease. Circulation. 2012;126(4):491–500.
14. Bernstein EF, Rhodes GA, Stuart SH, Coel MN, Fronk A. Toe pulse reappearance time in prediction of aortofemoral bypass success. Ann Surg. 1981;193(2):201–5.
15. Bertges DJ, Goodney PP, Zhao Y, Schanzer A, Nolan BW, Likosky DS, et al. The Vascular Study Group of New England Cardiac Risk Index (VSG-CRI) predicts cardiac complications more accurately than the Revised Cardiac Risk Index in vascular surgery patients. J Vasc Surg. 2010;52(3):674–83, 83 e1–83 e3.
16. Bhatt DL, Fox KA, Hacke W, Berger PB, Black HR, Boden WE, et al. Clopidogrel and aspirin versus aspirin alone for the prevention of atherothrombotic events. N Engl J Med. 2006;354(16):1706–17.
17. Blankenstein JD, de Jong SE, Prinsen M, van der Ham AC, Buth J, van Sterkenburg SM, et al. Two-year outcomes after conventional or endovascular repair of abdominal aortic aneurysms. N Engl J Med. 2005;352(23):2398–405.
18. Boccalon H, Lehert P, Mosnier M. [Effect of naftidrofuryl on physiological walking distance in patients with intermittent claudication]. Ann Cardiol Angeiol (Paris). 2001;50(3):175–82.
19. Bosch JL, Hunink MG. Meta-analysis of the results of percutaneous transluminal angioplasty and stent placement for aortoiliac occlusive disease. Radiology. 1997;204(1):87–96.

20. Bradbury AW, Adam DJ, Bell J, Forbes JF, Fowkes FG, Gillespie I, et al. Bypass versus Angioplasty in Severe Ischaemia of the Leg (BASIL) trial: A survival prediction model to facilitate clinical decision making. *J Vasc Surg*. 2010;51(5 Suppl):52S-68S.
21. Brighton TA, Eikelboom JW, Mann K, Mister R, Gallus A, Ockelford P, et al. Low-dose aspirin for preventing recurrent venous thromboembolism. *N Engl J Med*. 2012;367(21):1979–87.
22. Brooks B, Dean R, Patel S, Wu B, Molyneaux L, Yue DK. TBI or not TBI: that is the question. Is it better to measure toe pressure than ankle pressure in diabetic patients? *Diabet Med*. 2001;18(7):528–32.
23. Burger W, Chemnitius JM, Kneissl GD, Rucker G. Low-dose aspirin for secondary cardiovascular prevention – cardiovascular risks after its perioperative withdrawal versus bleeding risks with its continuation – review and meta-analysis. *J Intern Med*. 2005;257(5):399–414.
24. Cacoub PP, Bhatt DL, Steg PG, Topol EJ, Creager MA. Patients with peripheral arterial disease in the CHARISMA trial. *Eur Heart J*. 2009;30(2):192–201.
25. Canudas-Romo V. Three measures of longevity: time trends and record values. *Demography*. 2010;47(2):299–312.
26. Carbayo JA, Divison JA, Escribano J, Lopez-Abril J, Lopez de Coca E, Artigao LM, et al. Using ankle-brachial index to detect peripheral arterial disease: prevalence and associated risk factors in a random population sample. *Nutr Metab Cardiovasc Dis*. 2007;17(1):41–9.
27. Catalano M, Born G, Peto R. Prevention of serious vascular events by aspirin amongst patients with peripheral arterial disease: randomized, double-blind trial. *J Intern Med*. 2007;261(3):276–84.
28. Cleries R, Martinez JM, Valls J, Pareja L, Esteban L, Gispert R, et al. Life expectancy and age-period-cohort effects: analysis and projections of mortality in Spain between 1977 and 2016. *Public Health*. 2009;123(2):156–62.
29. Collins R, Burch J, Cranny G, Aguiar-Ibanez R, Craig D, Wright K, et al. Duplex ultrasonography, magnetic resonance angiography, and computed tomography angiography for diagnosis and assessment of symptomatic, lower limb peripheral arterial disease: systematic review. *BMJ*. 2007;334(7606):1257.
30. Connolly SJ, Eikelboom J, Joyner C, Diener HC, Hart R, Golitsyn S, et al. Apixaban in patients with atrial fibrillation. *N Engl J Med*. 2011;364(9):806–17.
31. Connolly SJ, Ezekowitz MD, Yusuf S, Eikelboom J, Oldgren J, Parekh A, et al. Dabigatran versus warfarin in patients with atrial fibrillation. *N Engl J Med*. 2009;361(12):1139–51.
32. Diehm C, Allenberg JR, Pittrow D, Mahn M, Tepohl G, Haberl RL, et al. Mortality and vascular morbidity in older adults with asymptomatic versus symptomatic peripheral artery disease. *Circulation*. 2009;120(21):2053–61.
33. Diehm C, Schuster A, Allenberg JR, Darius H, Haberl R, Lange S, et al. High prevalence of peripheral arterial disease and co-morbidity in 6880 primary care patients: cross-sectional study. *Atherosclerosis*. 2004;172(1):95–105.
34. Dieter R DR, Dieter RA III, ed. *Peripheral Aterial Disease*. . New York: McGraw Hill Companies; 2009.
35. Dörfler-Melly J KM, Prins MH , Büller HR . Antiplatelet and anticoagulant drugs for prevention of restenosis/reocclusion following peripheral endovascular treatment. *Cochrane Database Syst Rev* . 2005;( 1 ): CD002071. 2005.
36. Dormandy J, Heeck L, Vig S. The natural history of claudication: risk to life and limb. *Semin Vasc Surg*. 1999;12(2):123–37.
37. Doutremepuich C, Aguejouf O, Desplat V, Eizayaga FX. Aspirin therapy: an attempt to explain the events of prothrombotic complications after treatment discontinuation. *Thromb Haemost*. 2010;103(1):171–80.
38. Duda SH, Bosiers M, Lammer J, Scheiner D, Zeller T, Tielbeek A, et al. Sirolimus-eluting versus bare nitinol stent for obstructive superficial femoral artery disease: the SIROCCO II trial. *J Vasc Interv Radiol*. 2005;16(3):331–8.
39. Durazzo AE, Machado FS, Ikeoka DT, De Bernoche C, Monachini MC, Puech-Leao P, et al. Reduction in cardiovascular events after vascular surgery with atorvastatin: a randomized trial. *J Vasc Surg*. 2004;39(5):967–75; discussion 75–6.
40. Eugster T, Stierli P, Aeberhard P. Infrainguinal arterial reconstruction with autologous vein grafts: are the results for the in situ technique better than those of non-reversed bypass? A long-term follow-up study. *J Cardiovasc Surg (Torino)*. 2001;42(2):221–6.
41. Fallouh N, Chopra V. Statin withdrawal after major noncardiac surgery: risks, consequences, and preventative strategies. *J Hosp Med*. 2012;7(7):573–9.
42. Faulkner KW, House AK, Castleden WM. The effect of cessation of smoking on the accumulative survival rates of patients with symptomatic peripheral vascular disease. *Med J Aust*. 1983;1(5):217–9.
43. Feiring AJ, Wesolowski AA, Lade S. Primary stent-supported angioplasty for treatment of below-knee critical limb ischemia and severe claudication: early and one-year outcomes. *J Am Coll Cardiol*. 2004;44(12):2307–14.
44. Ferker BS, Spronk S, Colkesen EB, Hunink MG. Systematic review of guidelines on peripheral artery disease screening. *Am J Med*. 2012;125(2):198–208 e3.
45. Fleisher LA, Beckman JA, Brown KA, Calkins H, Chaikof E, Fleischmann KE, et al. ACC/AHA 2007 Guidelines on Perioperative Cardiovascular Evaluation and Care for Noncardiac Surgery: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 2002 Guidelines on Perioperative Cardiovascular Evaluation for Noncardiac Surgery): Developed in Collaboration With the American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Rhythm Society, Society of Cardiovascular Anesthesiologists, Society for Cardiovascular Angiography and Interventions, Society for Vascular Medicine and Biology, and Society for Vascular Surgery. *Circulation*. 2007;116(17):1971–96.
46. Fontaine R, Kim M, Kierny R. [Surgical treatment of peripheral circulation disorders]. *Helv Chir Acta*. 1954;21(5–6):499–533.
47. Fowkes FG, Murray GD, Butcher I, Heald CL, Lee RJ, Chambliss LE, et al. Ankle brachial index combined with Framingham Risk Score to predict cardiovascular events and mortality: a meta-analysis. *JAMA*. 2008;300(2):197–208.
48. Garimella PS, Hart PD, O'Hare A, DeLoach S, Herzog CA, Hirsch AT. Peripheral artery disease and CKD: a focus on peripheral artery disease as a critical component of CKD care. *Am J Kidney Dis*. 2012;60(4):641–54.
49. Greenhalgh RM, Brown LC, Powell JT, Thompson SG, Epstein D. Endovascular repair of aortic aneurysm in patients physically ineligible for open repair. *N Engl J Med*. 2010;362(20):1872–80.
50. Greenhalgh RM, Brown LC, Powell JT, Thompson SG, Epstein D, Sculpher MJ. Endovascular versus open repair of abdominal aortic aneurysm. *N Engl J Med*. 2010;362(20):1863–71.
51. Gruntzig AR. Percutaneous transluminal coronary angioplasty.

- Chronische arterielle Verschlusskrankheit der Extremitätenarterien. In: Angiologie, hrsg. von G. Heberer, G. Rau, W. Schoop, Thieme, Stuttgart. 1974:397.
53. Heintzen MP, Strauer BE. Peripheral vascular effects of beta-blockers. *Eur Heart J*. 1994;15 Suppl C:2–7.
  54. Hiatt WR. Medical treatment of peripheral arterial disease and claudication. *N Engl J Med*. 2001;344(21):1608–21.
  55. Hirsch AT, Criqui MH, Treat-Jacobson D, Regensteiner JG, Creager MA, Olin JW, et al. Peripheral arterial disease detection, awareness, and treatment in primary care. *JAMA*. 2001;286(11):1317–24.
  56. Hirsch AT, Haskal ZJ, Hertzler NR, Bakal CW, Creager MA, Halperin JL, et al. ACC/AHA 2005 Practice Guidelines for the management of patients with peripheral arterial disease (lower extremity, renal, mesenteric, and abdominal aortic): a collaborative report from the American Association for Vascular Surgery/Society for Vascular Surgery, Society for Cardiovascular Angiography and Interventions, Society for Vascular Medicine and Biology, Society of Interventional Radiology, and the ACC/AHA Task Force on Practice Guidelines (Writing Committee to Develop Guidelines for the Management of Patients With Peripheral Arterial Disease): endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation; National Heart, Lung, and Blood Institute; Society for Vascular Nursing; TransAtlantic Inter-Society Consensus; and Vascular Disease Foundation. *Circulation*. 2006;113(11):e463–654.
  57. Humphries MD, Pevec WC, Laird JR, Yeo KK, Hedayati N, Dawson DL. Early duplex scanning after infrainguinal endovascular therapy. *J Vasc Surg*. 2011;53(2):353–8.
  58. Huppert P, Tacke J, Lawall H. [S3 guidelines for diagnostics and treatment of peripheral arterial occlusive disease]. *Radiologe*. 2010;50(1):7–15.
  59. Imfeld S, Singer L, Degischer S, Aschwanden M, Thalhammer C, Labs KH, et al. Quality of life improvement after hospital-based rehabilitation or home-based physical training in intermittent claudication. *Vasa*. 2006;35(3):178–84.
  60. Jaeger KA, Amann-Vesti BR, Banyai M, Baumgartner I, Birrer M, Bounameaux M, et al. Schweizer Konsensus zur Therapie mit Thrombozyten-Funktionshemmern in der Angiologie. *Schweiz Med Forum*. 2009;9(39):690–3.
  61. Jonason T, Bergstrom R. Cessation of smoking in patients with intermittent claudication. Effects on the risk of peripheral vascular complications, myocardial infarction and mortality. *Acta Med Scand*. 1987;221(3):253–60.
  62. Koelmeij MJ, den Hartog D, Prins MH, Kromhout JG, Legemate DA, Jacobs MJ. Diagnosis of arterial disease of the lower extremities with duplex ultrasonography. *Br J Surg*. 1996;83(3):404–9.
  63. Korinth MC. Low-dose aspirin before intracranial surgery--results of a survey among neurosurgeons in Germany. *Acta Neurochir (Wien)*. 2006;148(11):1189–96; discussion 96.
  64. Korinth MC, Gilsbach JM, Weinzierl MR. Low-dose aspirin before spinal surgery: results of a survey among neurosurgeons in Germany. *Eur Spine J*. 2007;16(3):365–72.
  65. Kroger K, Stang A, Kondratieva J, Moebus S, Beck E, Schmermund A, et al. Prevalence of peripheral arterial disease – results of the Heinz Nixdorf recall study. *Eur J Epidemiol*. 2006;21(4):279–85.
  66. Lassila R, Lepantalo M. Cigarette smoking and the outcome after lower limb arterial surgery. *Acta Chir Scand*. 1988;154(11–12):635–40.
  67. Law M, Tang JL. An analysis of the effectiveness of interventions intended to help people stop smoking. *Arch Intern Med*. 1995;155(18):1933–41.
  68. Lederle FA, Johnson GR, Wilson SE, Littooy FN, Krupski WC, Bandyk D, et al. Yield of repeated screening for abdominal aortic aneurysm after a 4-year interval. *Aneurysm Detection and Management Veterans Affairs Cooperative Study Investigators*. *Arch Intern Med*. 2000;160(8):1117–21.
  69. Lehert P, Comte S, Gamand S, Brown TM. Naftidrofuryl in intermittent claudication: a retrospective analysis. *J Cardiovasc Pharmacol*. 1994;23 Suppl 3:S48–52.
  70. Macellari F, Paciaroni M, Agnelli G, Caso V. Perioperative stroke risk in nonvascular surgery. *Cerebrovasc Dis*. 2012;34(3):175–81.
  71. Makin AJ, Silverman SH, Lip GY. Antithrombotic therapy in peripheral vascular disease. *BMJ*. 2002;325(7372):1101–4.
  72. Mangano DT, Layug EL, Wallace A, Tateo I. Effect of atenolol on mortality and cardiovascular morbidity after noncardiac surgery. *Multicenter Study of Perioperative Ischemia Research Group*. *N Engl J Med*. 1996;335(23):1713–20.
  73. McFalls EO, Ward HB, Moritz TE, Goldman S, Krupski WC, Littooy F, et al. Coronary-artery revascularization before elective major vascular surgery. *N Engl J Med*. 2004;351(27):2795–804.
  74. McKenna M, Wolfson S, Kuller L. The ratio of ankle and arm arterial pressure as an independent predictor of mortality. *Atherosclerosis*. 1991;87(2–3):119–28.
  75. Melzer J, Brignoli R, Diehm C, Reichling J, Do DD, Saller R. Treating intermittent claudication with Tibetan medicine Padma 28: does it work? *Atherosclerosis*. 2006;189(1):39–46.
  76. Mills JL, Sr. Buerger's disease in the 21st century: diagnosis, clinical features, and therapy. *Semin Vasc Surg*. 2003;16(3):179–89.
  77. Murphy TP, Ariaratnam NS, Carney WI, Jr., Marcaccio EJ, Slaiby JM, Soares GM, et al. Aortoiliac insufficiency: long-term experience with stent placement for treatment. *Radiology*. 2004;231(1):243–9.
  78. Murphy TP, Cutlip DE, Regensteiner JG, Mohler ER, Cohen DJ, Reynolds MR, et al. Supervised exercise versus primary stenting for claudication resulting from aortoiliac peripheral artery disease: six-month outcomes from the claudication: exercise versus endoluminal revascularization (CLEVER) study. *Circulation*. 2012;125(1):130–9.
  79. Newman AB, Siscovick DS, Manolio TA, Polak J, Fried LP, Borhani NO, et al. Ankle-arm index as a marker of atherosclerosis in the Cardiovascular Health Study. *Cardiovascular Health Study (CHS) Collaborative Research Group*. *Circulation*. 1993;88(3):837–45.
  80. Norgren L, Hiatt WR, Dormandy JA, Nehler MR, Harris KA, Fowkes FG. Inter-Society Consensus for the Management of Peripheral Arterial Disease (TASC II). *J Vasc Surg*. 2007;45 Suppl S:S5–67.
  81. Norgren L, Hiatt WR, Harris KA, Lammer J. TASC II section F on revascularization in PAD. *J Endovasc Ther*. 2007;14(5):743–4.
  82. Patel MR, Mahaffey KW, Garg J, Pan G, Singer DE, Hacke W, et al. Rivaroxaban versus warfarin in nonvalvular atrial fibrillation. *N Engl J Med*. 2011;365(10):883–91.
  83. Poldermans D, Boersma E, Bax JJ, Thomson IR, Paelinck B, van de Ven LL, et al. Bisoprolol reduces cardiac death and myocardial infarction in high-risk patients as long as 2 years after successful major vascular surgery. *Eur Heart J*. 2001;22(15):1353–8.
  84. Poldermans D, Boersma E, Bax JJ, Thomson IR, van de Ven LL, Blankenstein JD, et al. The effect of bisoprolol on perioperative mortality and myocardial infarction in high-risk patients undergoing vascular surgery. Dutch

- Echocardiographic Cardiac Risk Evaluation Applying Stress Echocardiography Study Group. *N Engl J Med.* 1999;341(24):1789–94.
85. Poldermans D, Schouten O, Vidakovic R, Bax JJ, Thomson IR, Hoeks SE, et al. A clinical randomized trial to evaluate the safety of a noninvasive approach in high-risk patients undergoing major vascular surgery: the DECREASE-V Pilot Study. *J Am Coll Cardiol.* 2007;49(17):1763–9.
  86. Radack K, Deck C. Beta-adrenergic blocker therapy does not worsen intermittent claudication in subjects with peripheral arterial disease. A meta-analysis of randomized controlled trials. *Arch Intern Med.* 1991;151(9):1769–76.
  87. Rastan A, Noory E, Zeller T. Drug-eluting stents for treatment of focal infrapopliteal lesions. *Vasa.* 2012;41(2):90–5.
  88. Robertson L GM, Kovacs F. Drugs to prevent the re-occurrence for narrowing of blood vessels in peripheral arterial disease after the blood vessels have been surgically widened. *Cochrane Database Syst Rev*. Aug. 2012. 2012.
  89. Rodriguez LA, Cea-Soriano L, Martin-Merino E, Johansson S. Discontinuation of low dose aspirin and risk of myocardial infarction: case-control study in UK primary care. *BMJ.* 2011;343:d4094.
  90. Rooke TW, Hirsch AT, Misra S, Sidawy AN, Beckman JA, Findeiss LK, et al. 2011 ACCF/AHA focused update of the guideline for the management of patients with peripheral artery disease (updating the 2005 guideline): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines: developed in collaboration with the Society for Cardiovascular Angiography and Interventions, Society of Interventional Radiology, Society for Vascular Medicine, and Society for Vascular Surgery. *Catheter Cardiovasc Interv.* 2012;79(4):501–31.
  91. Ruffolo AJ, Romano M, Ciapponi A. Prostanoids for critical limb ischaemia. *Cochrane Database Syst Rev.* 2010(1):CD006544.
  92. Scheinert D, Katsanos K, Zeller T, Koppensteiner R, Commeau P, Bosiers M, et al. A prospective randomized multicenter comparison of balloon angioplasty and infrapopliteal stenting with the sirolimus-eluting stent in patients with ischemic peripheral arterial disease: 1-year results from the ACHILLES trial. *J Am Coll Cardiol.* 2012;60(22):2290–5.
  93. Schillinger M, Sabeti S, Loewe C, Dick P, Amighi J, Mlekusch W, et al. Balloon angioplasty versus implantation of nitinol stents in the superficial femoral artery. *N Engl J Med.* 2006;354(18):1879–88.
  94. Schlager O, Francesconi M, Haumer M, Dick P, Sabeti S, Amighi J, et al. Duplex sonography versus angiography for assessment of femoropopliteal arterial disease in a "real-world" setting. *J Endovasc Ther.* 2007;14(4):452–9.
  95. Schnyder-Joris C, Bonhomme F, Bonvini R, Fontana P. [Perioperative management of new antiplatelet drugs]. *Rev Med Suisse.* 2013;9(372):326–30.
  96. Spengel F, Clement D, Boccalon H, Liard F, Brown T, Lehert P. Findings of the Naftidrofuryl in Quality of Life (NIQOL) European study program. *Int Angiol.* 2002;21(1):20–7.
  97. Stewart KJ, Hiatt WR, Regensteiner JG, Hirsch AT. Exercise training for claudication. *N Engl J Med.* 2002;347(24):1941–51.
  98. Strandness DE, Jr., Dalman RL, Panian S, Rendell MS, Comp PC, Zhang P, et al. Effect of cilostazol in patients with intermittent claudication: a randomized, double-blind, placebo-controlled study. *Vasc Endovascular Surg.* 2002;36(2):83–91.
  99. Tendera M, Aboyans V, Bartelink ML, Baumgartner I, Clement D, Collet JP, et al. ESC Guidelines on the diagnosis and treatment of peripheral artery diseases: Document covering atherosclerotic disease of extracranial carotid and vertebral, mesenteric, renal, upper and lower extremity arteries: the Task Force on the Diagnosis and Treatment of Peripheral Artery Diseases of the European Society of Cardiology (ESC). *Eur Heart J.* 2011;32(22):2851–906.
  100. Tunis SR, Bass EB, Steinberg EP. The use of angioplasty, bypass surgery, and amputation in the management of peripheral vascular disease. *N Engl J Med.* 1991;325(8):556–62.
  101. Widmer M DF, Schmidli J. *Gefäßschirurgie in der Praxis.* Bern: Verlag Hans Huber, Hogrefe AG; 2009.
  102. Young PM, Mostardi PM, Glockner JF, Vrtiska TR, Macedo T, Haider CR, et al. Prospective Comparison of Cartesian Acquisition with Projection-like Reconstruction Magnetic Resonance Angiography with Computed Tomography Angiography for Evaluation of below-the-Knee Runoff. *J Vasc Interv Radiol.* 2013;24(3):392–9.
  103. Zeitler E, Schoop W, Zahnow W. The treatment of occlusive arterial disease by transluminal catheter angioplasty. *Radiology.* 1971;99(1):19–26.