

Genesung auf der Überholspur: Fast-Track-Chirurgie im DRG-Zeitalter

La guérison sur la voie de dépassement: Chirurgie fast-track à l'ère des DRG

Literatur / Références

- 1 Kehlet H,Wilmore DW. Multimodal strategies to improve surgical outcome. Am J Surg. 2002;183(6):630–41.
- 2 Adamina M, Kehlet H, Tomlinson GA, Senagore AJ, Delaney CP. Enhanced recovery pathways optimize health outcomes and resource utilization: a meta-analysis of randomized controlled trials in colorectal surgery. Surgery. 2011;149(6):830–40.
- 3 Kehlet H. Fast-track colorectal surgery. Lancet. 2008;371(9615):791–3.
- 4 Kehlet H,Wilmore DW. Evidence-based surgical care and the evolution of fast-track surgery. Ann Surg. 2008;248(2):189–98.
- 5 Wegmüller B,Bienlein M. Reshuffling the pack in the Swiss hospital market. World Hosp Health Serv. 2011;47(3):34–5.
- 6 Stephen AE,Berger DL. Shortened length of stay and hospital cost reduction with implementation of an accelerated clinical care pathway after elective colon resection. Surgery. 2003;133(3):277–82.
- 7 Roulin D, Donadini A, Gander S, Griesser AC, Blanc C, Hubner M, Schafer M, Demartines N. Cost-effectiveness of the implementation of an enhanced recovery protocol for colorectal surgery. Br J Surg. 2013;100(8):1108–14.
- 8 Longo WE, Virgo KS, Johnson FE, Oprian CA, Vernava AM, Wade TP, Phelan MA, Henderson WG, Daley J, Khuri SF. Risk factors for morbidity and mortality after colectomy for colon cancer. Dis Colon Rectum. 2000;43(1):83–91.
- 9 Schilling PL, Dimick JB, Birkmeyer JD. Prioritizing quality improvement in general surgery. J Am Coll Surg. 2008;207(5):698–704.
- 10 Ibrahim MS, Khan MA, Nizam I, Haddad FS. Peri-operative interventions producing better functional outcomes and enhanced recovery following total hip and knee arthroplasty: an evidence-based review. BMC Med. 2013;11:37.
- 11 Lemanu DP, Srinivasa S, Singh PP, Johannsen S, MacCormick AD, Hill AG. Optimizing perioperative care in bariatric surgery patients. Obes Surg. 2012;22(6):979–90.
- 12 Lu D, Wang X, Shi G. Perioperative enhanced recovery programmes for gynaecological cancer patients. Cochrane Database Syst Rev. 2012;12:CD008239.
- 13 Olsen MF,Wennberg E. Fast-track concepts in major open upper abdominal and thoracoabdominal surgery: a review. World J Surg. 2011;35(12):2586–93.
- 14 Zhu F, Lee A, Chee YE. Fast-track cardiac care for adult cardiac surgical patients. Cochrane Database Syst Rev. 2012;10:CD003587.
- 15 Delaney CP, Senagore AJ, Gerkin TM, Beard TL, Zingaro WM, Tomaszewski KJ, Walton LK, Poston SA. Association of surgical care practices with length of stay and use of clinical protocols after elective bowel resection: results of a national survey. Am J Surg. 2010;199(3):299–304.
- 16 Walter CJ, Smith A, Guillou P. Perceptions of the application of fast-track surgical principles by general surgeons. Ann R Coll Surg Engl. 2006;88(2):191–5.
- 17 Kube R, Ptak H, Steinert R, Sahm M, Schmidt U, Gastinger I, Lippert H. [Clinical value of laparoscopic surgery for colon cancer]. Chirurg. 2008;79(12):1145–50.
- 18 Berwick DM. Disseminating innovations in health care. JAMA. 2003;289(15):1969–75.
- 19 Delaney CP, Chang E, Senagore AJ, Broder M. Clinical outcomes and resource utilization associated with laparoscopic and open colectomy using a large national database. Ann Surg. 2008;247(5):819–24.
- 20 Raue W, Haase O, Junghans T, Scharfenberg M, Muller JM, Schwenk W. 'Fast-track' multimodal rehabilitation program improves outcome after laparoscopic sigmoidectomy: a controlled prospective evaluation. Surg Endosc. 2004;18(10):1463–8.
- 21 Jakobsen DH, Sonne E, Andreasen J, Kehlet H. Convalescence after colonic surgery with fast-track vs conventional care. Colorectal Dis. 2006;8(8):683–7.
- 22 DiFronzo LA, Yamin N, Patel K, O'Connell TX. Benefits of early feeding and early hospital discharge in elderly patients undergoing open colon resection. J Am Coll Surg. 2003;197(5):747–52.
- 23 Scharfenberg M, Raue W, Junghans T, Schwenk W. "Fast-track" rehabilitation after colonic surgery in elderly patients -is it feasible? Int J Colorectal Dis. 2007;22(12):1469–74.
- 24 Delaney CP, Fazio VW, Senagore AJ, Robinson B, Halverson AL, Remzi FH. 'Fast track' postoperative management protocol for patients with high co-morbidity undergoing complex abdominal and pelvic colorectal surgery. Br J Surg. 2001;88(11):1533–8.

- 25 Muller S, Zalunardo MP, Hubner M, Clavien PA, Demartines N. A fast-track program reduces complications and length of hospital stay after open colonic surgery. *Gastroenterology*. 2009;136(3):842–7.
- 26 Kariv Y, Delaney CP, Senagore AJ, Manilich EA, Hammel JP, Church JM, Ravas J, Fazio VW. Clinical outcomes and cost analysis of a "fast track" postoperative care pathway for ileal pouch-anal anastomosis: a case control study. *Dis Colon Rectum*. 2007;50(2):137–46.
- 27 Kehlet H, Dahl JB. Anaesthesia, surgery, and challenges in postoperative recovery. *Lancet*. 2003;362(9399):1921–8.
- 28 Smith I, Kranke P, Murat I, Smith A, O'Sullivan G, Soreide E, Spies C, in't VB. Perioperative fasting in adults and children: guidelines from the European Society of Anaesthesiology. *Eur J Anaesthesiol*. 2011;28(8):556–69.
- 29 Noblett SE, Watson DS, Huong H, Davison B, Hainsworth PJ, Horgan AF. Pre-operative oral carbohydrate loading in colorectal surgery: a randomized controlled trial. *Colorectal Dis*. 2006;8(7):563–9.
- 30 Lewis SJ, Andersen HK, Thomas S. Early enteral nutrition within 24 h of intestinal surgery versus later commencement of feeding: a systematic review and meta-analysis. *J Gastrointest Surg*. 2009;13(3):569–75.
- 31 Asgeirsson T, El-Badawi KI, Mahmood A, Barletta J, Luchtefeld M, Senagore AJ. Postoperative ileus: it costs more than you expect. *J Am Coll Surg*. 2010;210(2):228–31.
- 32 Muller SA, Rahbari NN, Schneider F, Warschkow R, Simon T, von FM, Bork U, Weitz J, Schmied BM, Buchler MW. Randomized clinical trial on the effect of coffee on postoperative ileus following elective colectomy. *Br J Surg*. 2012;99(11):1530–8.
- 33 Guenaga KF, Matos D, Wille-Jorgensen P. Mechanical bowel preparation for elective colorectal surgery. *Cochrane Database Syst Rev*. 2011;(9):CD001544.
- 34 Bundgaard-Nielsen M, Secher NH, Kehlet H. 'Liberal' vs. 'restrictive' perioperative fluid therapy--a critical assessment of the evidence. *Acta Anaesthesiol Scand*. 2009;53(7):843–51.
- 35 Chappell D, Jacob M, Hofmann-Kiefer K, Conzen P, Rehm M. A rational approach to perioperative fluid management. *Anesthesiology*. 2008;109(4):723–40.
- 36 Holte K, Sharrock NE, Kehlet H. Pathophysiology and clinical implications of perioperative fluid excess. *Br J Anaesth*. 2002;89(4):622–32.
- 37 Lowell JA, Schifferdecker C, Driscoll DF, Benotti PN, Bistrian BR. Postoperative fluid overload: not a benign problem. *Crit Care Med*. 1990;18(7):728–33.
- 38 Cheatham ML, Chapman WC, Key SP, Sawyers JL. A meta-analysis of selective versus routine nasogastric decompression after elective laparotomy. *Ann Surg*. 1995;221(5):469–76.
- 39 Nelson R, Edwards S, Tse B. Prophylactic nasogastric decompression after abdominal surgery. *Cochrane Database Syst Rev*. 2007;(3):CD004929.
- 40 Jottard K, Hoff C, Maessen J, van RB, van Berlo CL, Logeman F, Dejong CH. Life and death of the nasogastric tube in elective colonic surgery in the Netherlands. *Clin Nutr*. 2009;28(1):26–8.
- 41 Liu CL, Fan ST, Lo CM, Wong Y, Ng IO, Lam CM, Poon RT, Wong J. Abdominal drainage after hepatic resection is contraindicated in patients with chronic liver diseases. *Ann Surg*. 2004;239(2):194–201.
- 42 Petrowsky H, Demartines N, Rousson V, Clavien PA. Evidence-based value of prophylactic drainage in gastrointestinal surgery: a systematic review and meta-analyses. *Ann Surg*. 2004;240(6):1074–84.
- 43 Vlug MS, Wind J, Hollmann MW, Ubbink DT, Cense HA, Engel AF, Gerhards MF, van Wagensveld BA, van der Zaag ES, van Geloven AA, Sprangers MA, Cuesta MA, Bemelman WA. Laparoscopy in combination with fast track multimodal management is the best perioperative strategy in patients undergoing colonic surgery: a randomized clinical trial (LAFA-study). *Ann Surg*. 2011;254(6):868–75.
- 44 Adamina M, Gie O, Demartines N, Ris F. Contemporary perioperative care strategies. *Br J Surg*. 2013;100(1):38–54.
- 45 Kehlet H, Werner M, Perkins F. Balanced analgesia: what is it and what are its advantages in postoperative pain? *Drugs*. 1999;58(5):793–7.
- 46 Jin F, Chung F. Multimodal analgesia for postoperative pain control. *J Clin Anesth*. 2001;13(7):524–39.
- 47 McDonnell JG, O'Donnell B, Curley G, Heffernan A, Power C, Laffey JG. The analgesic efficacy of transversus abdominis plane block after abdominal surgery: a prospective randomized controlled trial. *Anesth Analg*. 2007;104(1):193–7.
- 48 Basse L, Werner M, Kehlet H. Is urinary drainage necessary during continuous epidural analgesia after colonic resection? *Reg Anesth Pain Med*. 2000;25(5):498–501.
- 49 Zaouter C, Kaneva P, Carli F. Less urinary tract infection by earlier removal of bladder catheter in surgical patients receiving thoracic epidural analgesia. *Reg Anesth Pain Med*. 2009;34(6):542–8.