

Warzen – Klinische Bilder und Therapie

Teil 2

Verrues – tableaux cliniques et traitement

Partie 2

Literatur/Références

1. Lin AN, Nakatsui T. Salicylic acid revisited. *International Journal of Dermatology* 1998;37:335–42.
2. Micali G, Dall'Oglio F, Nasca M, Tedeschi A. Treatment of cutaneous warts: an evidence-based approach. *American Journal of Clinical Dermatology* 2004;5:311–7.
3. Kwok CS, Gibbs S, Bennett C, Holland R, Abbott R. Topical treatments for cutaneous warts. *Cochrane Database Syst Rev* 2012 Sep 12;;9:CD001781. doi: 10.1002/14651858.CD001781.pub3.
4. Steele K, Shirodaria P, O'Hare M, Merrett JD, Irwin WG, Simpson DI. Monochloroacetic acid and 60% salicylic acid as a treatment of simple plantar warts: effectiveness and mode of action. *Br J Dermatol* 1988;118:537–43.
5. Burri P. Treatment of naevi and warts by topical chemotherapy with solcoderm. *Dermatologica* 1984;Suppl 1:52–7.
6. Hettich R. Solcoderm as a tool for the plastic surgeon. The treatment of verrucae. *Dermatologica* 1984;168 Suppl 1.
7. Yazar S, Basaran E. Efficacy of silver nitrate pencils in the treatment of common warts. *J Dermatol* 1994;21:329–33.
8. Schmidt H, Jacobsen F. Double-blind randomized clinical study on treatment of warts with fluorouracil-containing topical preparation [Ergebnis einer doppelblind und randomisiert durchgeführten klinischen Studie eines neuen Warzenmittels auf Basis Fluorouracil]. *Zeitschrift für Hautkrankheiten* 1981;56:41–3.
9. Wolff K. Result of a clinical trial of Verrumal (Hermal-Chemie) versus placebo; double-blind and randomised left/right study [Ergebnis einer klinischen Untersuchung von Verrumal gegen Placebo; doppelblind und randomisiert im Halbseitenversuch]. *Hermal internal report – Gutachten* 1980.

10. Iscimen A, Aydemir EH, Goksugur N, Engin B. Intralesional 5–fluorouracil, lidocaine and epinephrine mixture for the treatment of verrucae: a prospective placebo-controlled, single-blind randomized study. *J Eur Acad Dermatol Venereol* 2004;18:455–8.
11. Yazdanfar A, Farshchian M, Fereydoonejad M, Farshchian M. Treatment of common warts with an intralesional mixture of 5–fluorouracil, lidocaine, and epinephrine: a prospective placebo-controlled, double-blind randomized trial. *Dermatol Surg* 2008;34:656–9.
12. Batista CS, Atallah AN, Saconato H, da Silva EM. 5–FU for genital warts in non-immunocompromised individuals. *Cochrane Database Syst Rev* 2010;Apr 14;(4):CD006562. doi:10.1002/14651858.CD006562.pub2.
13. Weismann K, Kassis V. Treatment of condyloma acuminatum with 0,5% 5–fluorouracil-solutio. A double blind clinical trial. *Z Hautkr* 1982;57:810–6.
14. Perez AR WE, Piquero Martin J. Hypertonic saline solution vs intralesional bleomycin in the treatment of common warts. *Dermatology Venezuela* 1992;30:176–8.
15. Adalatkah H, Khalilollahi H, Amini N, Sadeghi-Bazargani H. Compared therapeutic efficacy between intralesional bleomycin and cryotherapy for common warts: A randomized clinical trial. *Dermatology Online Journal* 2007;13:4.
16. Dhar SB, Rashid MM, Islam A, Bhuiyan M. Intralesional bleomycin in the treatment of cutaneous warts: a randomized clinical trial comparing it with cryotherapy. *Indian Journal of Dermatology, Venereology and Leprology* 2009;75:262–7.
17. Epstein E. Persisting Raynaud’s phenomenon following intralesional bleomycin treatment of finger warts. *J Am Acad Dermatol* 1985;13:468–71.
18. Urbina González F, Cristobal Gil MC, Aguilar Martinez A, Guerra Rodriguez P, Sánchez de Paz F, García-Pérez A. Cutaneous toxicity of intralesional bleomycin administration in the treatment of periungual warts. *Arch Dermatol* 1986;122:974–5.
19. Herschthal J, McLeod MP, Zaiac M. Management of unguar warts. *Dermatol Ther* 2012;25:545–50.
20. Hellberg D, Svarrer T, Nilsson S, Valentin J. Self-treatment of female external genital warts with 0.5% podophyllotoxin cream (Condylone) versus weekly applications of 20% podophyllin solution. *Int J STD AIDS* 1995;6:257–61.
21. Syed TA, Lundin S, Ahmad SA. Topical 0.3% and 0.5% podophyllotoxin cream for self-treatment of condylomata acuminata in women. A placebo-controlled, double-blind study. *Dermatology* 1994;189:142–5.
22. Wang B, Wang B, Shao Y. A primary clinical trial of genital warts treated with domestic highly purified podophyllotoxin. *Chung Kuo I Hsueh Kuo Hsue Yuan Hsueh Pao* 1994;16:122–5.
23. Lacey CJ, Goodall RL, Tennvall GR, et al. Randomised controlled trial and economic evaluation of podophyllotoxin solution, podophyllotoxin cream, and podophyllin in the treatment of genital warts. *Sex Transm Infect* 2003;79:270–5.

24. Dawber RPR, Colver GB, Pringle F, Jackson A. Cutaneous cryosurgery: Principles and clinical practice. 2nd Edition. London: Taylor & Francis Ltd; 1997.
25. Bruggink SC, Gussekloo J, Berger MY, Zaaijer K, Assendelft WJ, de Waal MW. Cryotherapy with liquid nitrogen versus topical salicylic acid application for cutaneous warts in primary care: randomized controlled trial. *Canadian Medical Association Journal* 2010;182:1624–30.
26. Mi X, Chai W, Zheng H, Zuo YG, Li J. A randomized clinical comparative study of cryotherapy plus photodynamic therapy vs. cryotherapy in the treatment of multiple condylomata acuminata. *Photodermatol Photoimmunol Photomed* 2011;27:176–80.
27. Mulhem E, Pinelis S. Treatment of nongenital cutaneous warts. *Am Fam Physician* 2011;84:288–93.
28. Rüdlinger R. Moderne Warzenbehandlung. *Ther Umsch* 1998;55:498–501.
29. Mc Burney E, Rosen D. Carbon dioxide laser treatment of verrucae vulgaris. *J Dermatol Surg Oncol* 1984;10:45–8.
30. Mueller T, Carlson B, Lindy M. The use of carbon dioxide surgical laser for the treatment of verrucae. *J Am Podiatr Med Assoc* 1981;70:136–44.
31. Combemale P, Delolme H, Dupin M. Traitement de verrues. *Ann Dermatol Venereol* 1998;125:443–62.
32. Chen K, Chang B, Ju M, Zhang XH, Gu H. Comparative study of photodynamic therapy vs CO2 laser vaporization in treatment of condylomata acuminata: a randomized clinical trial. *Br J Dermatol* 2007;156:516–20.
33. Kauvar A, McDaniel D, Geronemus R. Pulsed dye laser treatment of warts. *Arch Fam Med* 1995;4:1035–40.
34. Kenton-Smith J, Tan S. Pulsed dye laser therapy for viral warts. *Br J Plast Surg* 1999;52:554–8.
35. Ross B, Levine V, Nehal K, Tse Y, Ashinoff R. Pulsed dye laser treatment of warts: an update. *Dermatol Surg* 1999;25:377–80.
36. Robson K, Cunningham N, Kruzan K, et al. Pulsed-dye laser versus conventional therapy in the treatment of warts: a prospective randomized trial. *J Am Acad Dermatol* 2000.;43:275–80.
37. Passeron T, Sebban K, Mantoux F, Fontas E, Lacour JP, Ortonne JP. 595 nm pulse dye laser therapy for viral warts: a single-blind randomized comparative study versus placebo. *Ann Dermatol Venereol* 2007;134:135–9.
38. Stender I, Lock Andersen J, Wulf H. Recalcitrant hand and foot warts successfully treated with photodynamic therapy with topical 5-aminolaevulinic acid: a pilot study. *Clin Exp Dermatol* 1999;24:154–9.

39. Stender I, Na R, Fogh H, Gluud C, Wulf H. Photodynamic therapy with 5-aminolaevulinic acid or placebo for recalcitrant foot and hand warts: randomised double blind trial. *Lancet* 2000;355:963–6.
40. Fuchs SM, Fluhr JW, Bankova L, Tittelbach J, Hoffmann G, Elsner P. Photodynamic therapy (PDT) and waterfiltered infrared A (wIRA) in participants with recalcitrant common hand and foot warts. *GMS German Medical Science* 2004;2:Doc08.
41. Ying Z, Li X, Dang H. 5-aminolevulinic acid-based photodynamic therapy for the treatment of condylomata acuminata in Chinese patients: a meta-analysis. *Photodermatol Photoimmunol Photomed* 2013;29:149–59.
42. Liang J, Lu XN, Tang H, Zhang Z, Fan J, Xu JH. Evaluation of photodynamic therapy using topical aminolevulinic acid hydrochloride in the treatment of condylomata acuminata: a comparative, randomized clinical trial. *Photodermatol Photoimmunol Photomed* 2009;25:293–7.
43. Hoffmann G. Principles and working mechanisms of water-filtered infrared-A (wIRA) in relation to wound healing. *GMS Krankenhhyg Interdiszip* 2007;2:Doc54.
44. Eriksen K. Treatment of the common wart by induced allergic inflammation. *Dermatologica* 1980;160:161–6.
45. Claudy AL, Roche H, Gogue Y. Immunotherapy of multiple recurring warts. Reassessment of the use of dinitrochlorobenzene. *Ann Dermatol Venereol* 1980;107:551–3.
46. Sanders BB, Smith KW. Dinitrochlorobenzene immunotherapy of human warts. *Cutis* 1981 27:389–92.
47. Lee S, Cho CK, Kim JG, Chun SI. Therapeutic effect of dinitrochlorobenzene (DNCB) on verruca plana and verruca vulgaris. *Int J Dermatol* 1984;23:624–6.
48. Rosado-Cancino MA, Ruiz-Maldonado R, Tamayo L, Laterza AM. Treatment of multiple and stubborn warts in children with 1-chloro-2,4-dinitrobenzene (DNCB) and placebo *Dermatologia Revista Mexicana* 1989;33:245–52.
49. Wilson P. Immunotherapy v cryotherapy for hand warts; a controlled trial (abstract). *Scottish Medical Journal* 1983;28:191.
50. Buckley DA, Keane FM, Munn SE, Fuller LC, Higgins EM, Du Vivier AW. Recalcitrant viral warts treated by diphenycprone immunotherapy. *Br J Dermatol* 1999;141:292–6.
51. Choi Y, Kim do H, Jin SY, Lee AY, Lee SH. Topical immunotherapy with diphenylcyclopropenone is effective and preferred in the treatment of periungual warts. *Ann Dermatol Venereol* 2013;25:434–9.
52. Choi MH, Seo SH, Kim IH, Son SW. Comparative study on the sustained efficacy of diphenycprone immunotherapy versus cryotherapy in viral warts. *Pediatr Dermatol* 2008;25:398–9.

53. Pires MC, Martins JM, Montealegre F, Gatti FR. Vitiligo after diphencyprone for alopecia areata. *Dermatol Res Pract* 2010;171265. doi: 10.1155/2010/. Epub 2010 May 11.
54. Horn TD, Johnson SM, Helm RM, Roberson PK. Intralesional immunotherapy of warts with mumps, candida and trichophyton skin test antigens. *Archives of Dermatology* 2005;141:589–94.
55. Dahl M. Imiquimod: an immune response modifier *J Am Acad Dermatol* 2000;43:S1–5.
56. Edwards L. Imiquimod in clinical practice. *J Am Acad Dermatol* 2000;43:S12–7.
57. Edwards L, Ferenczy A, Eron L, et al. Self-administered topical 5% imiquimod cream for external anogenital warts. *Arch Dermatol* 1998;134:25–30.
58. Diamantis ML, Bartlett BL, Tying SK. Safety, efficacy & recurrence rates of imiquimod cream 5% for treatment of anogenital warts. *Skin Therapy Letters* 2009;14:1–3, 5.
59. Aldara 3M 2000 a. Vehicle-controlled dose ranging study of Aldara 5% cream for the treatment of plantar warts. 1319IMI. (Short report of an RCT supplied by 3M in 2000). Data on file. 19 May 2000.
60. Aldara 3M 2000 b. Vehicle-controlled dose ranging study of Aldara 5% cream. 1321IMI. (Short report of an RCT supplied by 3M in 2000). Data on file. 14 April 2000.
61. Stockfleth E, Meyer T. The use of sinecatechins (polyphenon E) ointment for treatment of external genital warts. *Expert Opin Biol Ther* 2012;12:783–93.
62. Tatti S, Stockfleth E, Beutner KR, et al. Polyphenon E: a new treatment for external anogenital warts. *Br J Dermatol* 2010;162:176–84.
63. Stockfleth E, Beti H, Orasan R, et al. Topical Polyphenon E in the treatment of external genital and perianal warts: a randomized controlled trial. *Br J Dermatol* 2008;158:1329–38.
64. Gross G, Meyer KG, Pres H, Thielert C, Tawfik H, Mescheder A. A randomized, double-blind, four-arm parallel-group, placebo-controlled Phase II/III study to investigate the clinical efficacy of two galenic formulations of Polyphenon E in the treatment of external genital warts. *J Eur Acad Dermatol Venereol* 2007;21:1404–12.
65. Davis M, Gostout B, Mc Govern R, Persing D, Schut R, Pittelkow M. Large plantar wart caused by human papillomavirus-66 and resolution by topical codofovir therapy. *J Am Acad Dermatol* 2000;43:340–3.
66. Wenner R, Askari SK, Cham PMH, Kedrowski DA, Liu A, Warshaw EM. Duct tape for the treatment of common warts in adults: A double-blind randomized controlled trial. *Arch Dermatol* 2007;143:309–13.
67. de Haen M, Spigt MG, van Uden CJT, van Neer P, Feron FJM, Knottnerus A. Efficacy of duct tape vs placebo in the treatment of verruca vulgaris (warts) in primary school children. *Arch Pediatr Adolesc Med* 2006;160:1121–5.

68. Kim SY, Jung SK, Lee SG, Yi SM, Kim JH, Kim IH. New alternative combination therapy for recalcitrant common warts: the efficacy of imiquimod 5% cream and duct tape combination therapy. *Ann Dermatol* 2013;25:261–3.
69. Joseph R, Pulimood SA, Abraham P, John GT. Successful treatment of verruca vulgaris with Thuja occidentalis in a renal allograft recipient. *Indian J Nephrol* 2013;23:362–4.
70. Khan MT, Cerio R, Watt R, Khan MTSnr. A double-blind placebo study of topical thuja occidentalis on verruca pedis in children and adults. *J Eur Acad Dermatol Venereol* 1999;12:S251–2 (P-388).
71. Khan MT, Cerio R, Watt R, Khan MT. Phytotherapy with subfractions of topical Thuja Occ on verruca pedis. *Br J Dermatol* 2000;143:(Suppl 57):33.
72. Zedan H, Hofny ER, Ismail SA. Propolis as an alternative treatment for cutaneous warts. *Int J Dermatol* 2009;48:1246–9.
73. Millar BC, Moore JE. Successful topical treatment of hand warts in a paediatric patient with tea tree oil (*Melaleuca alternifolia*). *Complement Ther Clin Pract* 2008;14:225–7.
74. Bhushan M, Beck MH. Allergic contact dermatitis from tea tree oil in a wart patient. *Contact Dermatitis* 1997;36:117–8.
75. Pirker C, Hausen BM, Uter W, et al. Sensitization to tea tree oil in Germany and Austria. A multicenter study of the German Contact Dermatitis Group. *J Dtsch Dermatol Ges* 2003;1:629–34.
76. Labrecque M, Audet D, Latulippe L, Drouin J. Homeopathic treatment of plantar warts. *Can Med Assoc* 1992;146:1749–53.
77. Kainz J, Kozel G, Haidvogel M, Smolle J. Homeopathic versus placebo therapy of children with warts on the hands: a randomized, double-blind clinical trial. *Dermatology* 1996;193:318–20.
78. Meineke V, Reichrath J, Reinhold U, Tilgen W. Verrucae vulgares in children: successful simulated X-ray treatment (a suggestion-based therapy). *Dermatology* 2002;204:287–9.
79. Spanos N, Stentstrom R, Johnston J. Hypnosis, placebo and suggestion in the treatment of warts. *Psychosom Med* 1988;50:245–60.
80. Spanos NP, Williams V, Gwynn MI. Effects of hypnotic, placebo, and salicylic acid treatments on wart regression. *Psychosom Med* 1990;52:109–14.
81. Harkness E, Abbot N, Ernst E. A randomized trial of distant healing for skin wart. *Am J Med* 2000;15:507–8.
82. Gillison ML, Chaturvedi AK, Lowy DR. HPV prophylactic vaccines and the potential prevention of noncervical cancers in both men and women. *Cancer* 2008;113 (10 Suppl):3036–46.

83. Read TR, Hocking JS, Chen MY, Donovan B, Bradshaw CS, Fairley CK. The near disappearance of genital warts in young women 4 years after commencing a national human papillomavirus (HPV) vaccination programme. *Sex Transm Infect* 2011;87:544–7.