

## Flüssigkeits- und Volumenersatz in der Intensivmedizin. Ein Update über den aktuellen Stand des Irrtums

### Substitution hydrique et volumique en Médecine intensive. Update sur l'état actuel de l'erreur

#### Literatur / Références

- 1 Churton E. Leeds General Infirmary: A case of scirrus of the pylorus, with excessive vomiting; repeated intravenous injection of saline solution; remarks. *Lancet* 1888;132:620–621
- 2 Chowdhury AC, Cox EF, Francis ST, Lobo DN. A randomized, controlled, double blind crossover study on the effects of 2 L infusions of 0.9% saline and plasma lyte 148 on renal blood flow velocity and renal cortical tissue perfusion in healthy volunteers. *Ann Surg* 2012;256:18–24
- 3 Nor'azim MY, Bellomo R, Hegarty C, Story D, Ho L, Bailey M. Association Between a Chloride-Liberal vs Chloride-Restrictive Intravenous Fluid Administration Strategy and Kidney Injury in Critically Ill Adults. *JAMA*. 2012;308:1566–72
- 4 Shaw AD, Bagshaw SM, Goldstein SL et al. Major complications, mortality, and resource utilization after open abdominal surgery: 0.9% saline compared to Plasma-Lyte. *Ann Surg* 2012;255:821–9.
- 5 Helfen T. Der exakte Volumeneffekt der Vollelektrolytlösung Ringer-Laktat unter den Bedingungen der akuten normovolämischen Hämodilution. Dissertation zum Erwerb des Doktorgrades der Medizin an der Medizinischen Fakultät der Ludwig-Maximilians-Universität zu München. 2012
- 6 Shoemaker WC. Comparison of the relative effectiveness of whole blood transfusions and various types of fluid therapy in resuscitation. *Crit Care Med*, 1976; 4:71,
- 7 Haljame H. Rational for the use of colloids in the treatment of shock and hypovolemia. *Acta Physiol Scand*, 1985; 29 (suppl. 82)
- 8 Schott U, Lindbom LO, Sjöstrand U. Hemodynamic effects of colloid concentration in experimental hemorrhage: A comparison of Ringers acetate, 3% dextran-60, and 6% dextran 70. *Crit Care Med*, 1988; 16: 346
- 9 Hauser CJ, Shoemaker WC, Turpin I, Goldberg SJ. Oxygen transport responses to colloids and crystalloids in critically ill surgical patients. *Surg Gynecol Obstet*, 1980; 150: 811
- 10 Kasotakis G, Sideris A, Yuchiao Yang Y, et al. Aggressive early crystalloid resuscitation adversely affects outcomes in adult blunt trauma patients: An analysis of the Glue Grant database *J Trauma Acute Care Surg* 2013;74: 1215–22
- 11 Neff TA, Doelberg M, Jungheinrich C, Sauerland A, Spahn DR, Stocker R. Repetitive large-dose infusion of the novel hydroxyethyl starch 130/0.4 in patients with severe head injury. *Anesth Analg* 2003;96:1453–9
- 12 Béchir M, Puhan M, Neff SB, Guggenheim M, Wedler V, Stover JF, Stocker R, et al. Early fluid resuscitation with hyperoncotic hydroxyethyl starch 200/0.5 (10%) in severe burn injury. *Crit Care*. 2010;14:R123
- 13 Morard JC, Rutishauser E, Chatillon JJ. Anatomoclinical demonstration of 5 cases of osmotic nephrosis *Urol Medicae Chir.* 1955;61:612–25
- 14 Dickenmann M, Oettl T, Mihatsch MJ. Osmotic nephrosis: Acute kidney injury with accumulation of proximal tubular lysosomes due to administration of exogenous solutes. *American Journal of Kidney Diseases* 2008; 51:491–503
- 15 Rozich JD, Paul RV. Acute renal failure precipitated by elevated colloid osmotic pressure. *Am J Med* 1989; 87:359–60
- 16 Brunkhorst FM, Engel Ch, Frank Bloos F, Meier-Hellmann F, Ragaller M, Weiler N. et al. Intensive Insulin Therapy and Pentastarch Resuscitation in Severe Sepsis. *N Engl J Med* 2008; 358:125–39
- 17 James MF1, Michell WL, Joubert IA, Nicol AJ, Navsaria PH, Gillespie RS Resuscitation with hydroxyethyl starch improves renal function and lactate clearance in penetrating trauma in a randomized controlled study: the FIRST trial (Fluids in Resuscitation of Severe Trauma). *Br J Anaesth*. 2011;107:693–702
- 18 Guidet B, Martinet O, Boulain T, Philippart F, Poussel JF, Maizel J, Forceville X. Assessment of hemodynamic efficacy and safety of 6% hydroxyethylstarch 130/0.4 vs. 0.9% NaCl fluid replacement in patients with severe sepsis: The CRYSTMAS study. *Crit Care* 2012, 16:R94
- 19 Perner A, Haase N, Guttormsen AB, Tenhunen J, Klemenzson G, Åneman A, et al. Hydroxyethylstarch 130/0.42 versus Ringer's acetate in severe sepsis. *N Engl J Med* 2012;367:124–34
- 20 Myburgh JA, Finfer S, Bellomo R, Billot L, Cass A, Gattas D, Glass P, et al. Hydroxyethyl starch or saline for fluid resuscitation in intensive care. *N Engl J Med* 2012;367:1901–11
- 21 Annane D, Siami S, Jaber S, Martin C, Elatrous S, Descorps Declère A, et al. Effects of Fluid Resuscitation With Colloids vs Crystalloids on Mortality in Critically Ill Patients Presenting With Hypovolemic Shock. The CRISTAL Randomized Trial. *JAMA*. 2013;310:1809–17
- 22 The SAFE Study Investigators A Comparison of Albumin and Saline for Fluid Resuscitation in the Intensive Care Unit *N Engl J Med* 2004; 350:2247–56

23 Caironi P, Tognoni G, Masson S, et al. for the ALBIOS Study Investigators. Albumin Replacement in Patients with Severe Sepsis or Septic Shock *N Engl J Med* 2014; 370:1412–1421.