

ARDS: ventilate gently, relax, dry and prone

Literatur / Références

- 1 Piehl MA, Brown RS. Use of extreme position changes in acute respiratory failure. Crit Care Med 1976;4:13–4.
- 2 Gattinoni L, Tognoni G, Pesenti A, Taccone P, Mascheroni D, Labarta V et al. Effect of prone positioning on the survival of patients with acute respiratory failure. N Eng J Med 2001;345:568–73.
- 3 Guerin C, Gaillard S, Lemasson S, Ayzac L, Girard R, Beuret P et al. Effects of systematic prone positioning in the hypoxemic acute respiratory failure: a randomized controlled trial. JAMA 2004;292:2379–87.
- 4 Mancebo J, Fernández R, Blanch L, Rialp G, Gordo F, Ferrer M et al. A multicenter trial of prolonged prone ventilation in severe acute respiratory distress syndrome. Am J Respir Crit Care Med 2006;173:1233–9.
- 5 Taccone P, Pesenti A, Latini R, Polli F, Vagginelli F, Mietto C et al. Prone positioning in patients with moderate and severe acute respiratory distress syndrome: a randomized controlled trial. JAMA 2009;302:1977–84.
- 6 Guerin C, Reignier J, Richard JC, Beuret P, Gacouin A, Boulain T et al. Prone positioning in severe acute respiratory distress syndrome. N Eng J Med 2013; 368(23):2159–68.
- 7 Ferguson ND, Cook DJ, Guyatt GH, Mehta S, Hand L, Austin P et al. High-frequency oscillation in early acute respiratory distress syndrome. N Eng J Med 2013;368(9):795–805.
- 8 Schmidt M, Zogheib E, Rozé H, Repesse X, Lebreton G, Luyt CE et al. The PRESERVE mortality risk score and analysis of long-term outcomes after extracorporeal membrane oxygenation for severe acute respiratory distress syndrome. Intensive Care Med 2013;39(10):1704–13.